arXiv:1304.1915v2 [math.CV] 24 May 2013

CONFORMAL MAPS AND UNCONTROLLABLY WIGGLY BOUNDARIES

TIMOTHY H. MCNICHOLL

Abstract. It is shown that there is a computable conformal map of the unit disk onto a domain D that has a computable boundary extension even though the boundary of D is not effectively locally connected. The proof encodes an arbitrary *c.e.* set into the local connectivity of the boundary of D.

1. Introduction

Our goal here is to investigate the information necessary to compute the boundary extension of a conformal map. In [21], it is shown that local connectivity information for the boundary of the range is sufficient; that is the ability to predict an upper bound on the diameter of paths in the boundary between nearby points. Here, we will show that local connectivity information is not *necessary.* Namely, we show that there is a computable and conformal map ϕ of the unit disk (that is, the open disk with center 0 and radius 1) onto a domain D so that ϕ has a computable boundary extension but the boundary of D is not effectively locally connected. That is, no algorithm can predict an upper bound on the diameter of paths in the boundary between nearby points.

We will build D by encoding an arbitrary *c.e.* set into the local connectivity of its boundary. The boundary of this domain D is essentially a polygon with infinitely many vertices. So, we will use the technology developed by Andreev and McNicholl in [1] to create a representation of ϕ in terms of the functions of potential theory. We then use this representation of ϕ to show that its boundary extension is computable.

We give background from complex analysis and computability in Section 2. In Section 3, we prove a new estimate on the boundary behavior of the inverse of a conformal map. Then, in Section 4 we construct the domain D and its boundary and prove that the boundary of D is not effectively locally connected. In Section 5 we build a representation of the map ϕ in terms of the functions of potential theory. We prove the computability of the constituents of this representation in Section 6. We demonstrate the computability of the boundary extension of ϕ in Section 7. Concluding observations and further questions for investigation are discussed in Section 8.

2. Background

2.1. **Background from topology and complex analysis.** Let $\mathbb{C}$ denote the complex plane. A *domain* is a subset of $\mathbb{C}$ that is open and connected. A domain

---

1991 *Mathematics Subject Classification.* 30C30, 30E10, 03D78, 03F60, 54D05.

Key words and phrases. boundary behavior of conformal maps, approximation, computational complex analysis, computable analysis, effective local connectivity.

D is *simply connected* if it has the property that whenever a Jordan curve γ is included in D its interior is as well.

Let $\mathbb{D}$ denote the *unit disk*; that is the open disk whose center is the origin and whose radius is 1. The boundary of $\mathbb{D}$ will be called the *unit circle*. The Riemann Mapping Theorem states that if D is a simply connected domain with at least two boundary points, then there is a conformal map of the unit disk onto D. If D is also a Jordan domain (ie the boundary of D is a Jordan curve), then the Carathéodory Theorem states there is a homeomorphic extension of ϕ to the closure of the unit disk.

A topological space X is *locally connected* if for every point $p \in X$ and every neighborhood of p, U, there is a connected neighborhood of p, V, that is included in U. If X is a compact metric space, then X is effectively locally connected if and only if it is *uniformly locally arcwise connected*; that is, for every $\epsilon > 0$, there is a $\delta > 0$ so that whenever $p, q \in X$ are distinct, if $d(p, q) < \delta$, then there is an arc in X from p to q whose diameter is smaller than ϵ. The Hahn-Mazurkiewicz Theorem states that a Hausdorff topological space X is the range of a continuous function on $[0, 1]$ if and only if it is compact, metrizable, connected, and locally connected.

If ϕ is a conformal map of the unit disk onto a bounded domain D, and if the boundary of D is not a Jordan curve but is at least locally connected, then ϕ has a continuous extension to the closure of the unit disk. This extension is called the *boundary extension* of ϕ and is denoted by ϕ as well. By the converse of the Hahn-Mazurkiewicz Theorem, if a conformal map of the unit disk onto a domain D has a boundary extension, then the boundary of D is locally connected. Boundary extensions of conformal maps play an important role in the solution of Dirichlét problems.

In the subsequent sections, a number of results from the theory of analytic and harmonic functions are quoted without proofs. Except where noted, demonstrations can be found in Garnett and Marshall [12], Nehari [25], and Pommerenke [26].

2.2. **Background from computable analysis.** With regards to computability over the natural numbers, in particular the meaning of the term *computable* as it applies to sets of natural numbers and to functions that map natural numbers to natural numbers, and also in particular to the meaning of *computably enumerable*, we refer the reader to standard texts such as Cutland [9], and Cooper [7]. A common example of a computably enumerable set that is incomputable is *Turing's Halting Set*, which can be defined as the set of all codes of Turing machines that halt on at least one input.

With regards to computability over continuous spaces such as the complex plane, there is no standard text to which we can refer the reader. Two important works are the Banach space approach of Pour-El and Richards [27] and the Type-Two Effectivity approach in Weihrauch [28]. The latter establishes a computability theory on any second-countable T_0 space. There is also the point-free approach of Kalantari and Welch [17]. On all spaces common to their purviews, these three approaches are equivalent. Complexity of computation with real numbers is treated in Ko [19]. See also Cook and Kawamura [18]. Here, we only consider computability on the complex plane and its hyperspaces. So, we will use the following definitions which are equivalent in this setting to these other frameworks.

To begin, a *rational interval* is an interval whose endpoints are rational numbers. An *open rational rectangle* is a cartesian product of two open rational intervals. A *closed rational rectangle* is a cartesian product of two closed rational intervals.

A point $p \in \mathbb{C}$ is *computable* if there is an algorithm that, given any natural number k as input, computes a rational point $q \in \mathbb{C}$ such that $|q - p| < 2^{-k}$. That is, it is possible to compute arbitrarily good approximations of p. All algebraic numbers are computable as are many transcendental numbers such as π and e. However, most points are incomputable. In particular, if A is an incomputable set of natural numbers, then

$$\sum_{n \in A} 2^{-n}$$

is an incomputable real.

A sequence $\{z_n\}_{n=0}^{\infty}$ is *computable* if there is an algorithm that, given any $k, n \in \mathbb{N}$ as input, computes a rational point $q \in \mathbb{C}$ such that $|q - z_n| < 2^{-k}$. So, if c is a computable real, and if $z_n = c$ for all n, then $\{z_n\}_{n \in \mathbb{N}}$ is computable. On the other hand, if A is an incomputable set of natural numbers, and if

$$z_n = \begin{cases} 1 & \text{if } n \in A \\ 0 & \text{if } n \notin A \end{cases}$$

then $\{z_n\}_{n \in A}$ is incomputable.

Suppose f is a function that maps complex numbers to complex numbers. We say that f is *computable* if there is an algorithm M that meets the following three criteria.

- **Approximation:** Given an open rational rectangle R_1 as input, either M does not halt, or it produces a rational rectangle R_2 as output.
- **Correctness:** If M halts on a rational rectangle R_1, and if f is defined at $z \in R_1$, then $f(z) \in R_2$.
- **Convergence**: If U is a neighborhood of $f(z)$, and if R is a rational rectangle that contains z, then on a rational rectangle $R_1 \subseteq R$ that contains z, M halts and computes a rational rectangle that is contained in U.

In other words, f is computable if for each z in its domain we can compute arbitrarily good approximations of $f(z)$ from sufficiently good approximations of z. Most functions that occur in practice are computable. In addition, all computable functions are continuous. However, if z_0 is an incomputable point, then $f(z) = z_0$ gives a continuous function that is incomputable.

An open subset of the plane U is *computable* if the set of closed rational rectangles included in U is computably enumerable. So, for example, if r is a computable positive real, and if z_0 is a computable point then $D_r(z_0)$ is a computable open set.

A closed subset of the plane C is *computable* if the set of open rational rectangles that contain a point of C is computably enumerable. For example, it follows from the definitions that if $f : [0,1] \to \mathbb{C}$ is computable, then its range is a computable closed set.

A compact subset of the plane C is *computably compact* if there is an algorithm that, given a natural number k as input, computes a finite set of rational rectangles $\{R_1, \ldots, R_n\}$ with the following properties.

- The diameter of each R_j is smaller than 2^{-k}.
- Each R_j contains a point of C.
- $C \subseteq \bigcup_j R_j$.

It follows that a computably compact set is a computably closed set [5]. Intuitively, computable compactness corresponds to the ability to plot the set on a computer screen with arbitrarily good resolution. If $f : [0,1] \to \mathbb{C}$ is computable, then it follows from the definitions that its range is computably compact.

We say that $X \subseteq \mathbb{C}$ is *effectively locally connected* if there is a computable function $g : \mathbb{N} \to \mathbb{N}$ with the property that whenever $k \in \mathbb{N}$ and p, q are distinct points of X, if $|p - q| \leq 2^{-g(k)}$, then X contains an arc from p to q whose diameter is smaller than 2^{-k}. A definition of "effectively locally connected" that is a faithful translation of the standard definition of "locally connected" into the framework of Type Two Effectivity appears in Brattka [4]. In Daniel and McNicholl [10], it is shown that for computably compact subsets of the plane, this notion is equivalent to the one just given. Effective local connectivity is a way of controlling the amount of zigzagging in the boundary of a domain which then allows for the construction of accessing arcs. That is, arcs from that proceed from a point in the domain to a point on the boundary without intersecting any other point of the boundary. The ability to compute such arcs is used to compute boundary extensions of conformal maps in McNicholl [22]. Computable aspects of accessing arcs are explored in McNicholl [23].

Suppose f is a function that maps complex numbers to complex numbers. A function $g : \mathbb{N} \to \mathbb{N}$ is a *modulus of continuity* for f if $|f(z_1) - f(z_2)| < 2^{-k}$ whenever $k \in \mathbb{N}$ and z_1, z_2 are points in the domain of f such that $|z_1 - z_2| \leq 2^{-g(k)}$. It is well-known that a computable function $f : [0,1] \to \mathbb{C}$ has a computable modulus of continuity. It is also well-known that the range of such a function is computably compact.

In 1912, P. Koebe [20] outlined a constructive proof of the Riemann Mapping Theorem (see also Henrici [15]). A proof of the Riemann Mapping Theorem in the framework of Bishop that builds on Koebe's ideas is developed in Cheng [6] and in Bishop and Bridges [3]. An effective Riemann Mapping Theorem appears in Hertling [16]. The essence of this result is that if a domain D is simply connected and has at least two boundary points, and if it is also the case that D and its boundary can be computed (in a sense made precise in Section 2), then it is possible to compute a conformal map ϕ of the unit disk onto D. Furthermore, the proof of this result is *uniform* in that it yields an algorithm that transforms algorithms for computing D and its boundary into an algorithm for computing ϕ. It is also shown in the same paper that if a conformal map of the unit disk onto D can be computed, then D and its boundary can be computed as well. Again, the proof is uniform. It is observed in Pour-El and Richards [27] that it is possible to compute a domain D whose boundary is incomputable. Together, these results pin down D and its boundary as the information necessary to compute ϕ. The complexity of computing ϕ when D is a Jordan domain is explored in Binder, Braverman, and Yampolsky [2].

The information-theoretic complexity of computing boundary extensions of conformal maps onto simply-connected Jordan domains was determined in McNicholl [22]. Namely, it is shown that a computable conformal map of the unit disk onto a Jordan domain D has a computable boundary extension if and only if the boundary of D is effectively locally connected. Again, the proof is uniform.

These developments lead to the investigation of the information required to compute the boundary extension of a conformal map onto a non-Jordan domain whose boundary is locally connected. In McNicholl [21], it is shown that a boundary extension can be computed if the domain D and its boundary are computable and its boundary is effectively locally connected. So, at first glance, it seems that effective local connectivity provides the right amount of information necessary for computing boundary extensions. However, Couch, Daniel, and McNicholl observed in [8], based on work of Miller [24] and Gu, Lutz, and Mayordomo [14], that there is a subset of the plane that is not effectively locally connected even though it forms the range of a computable map on $[0, 1]$. Here we strengthen this result by showing that it is possible to compute a conformal map ϕ of the unit disk onto a domain D in such a way that ϕ also has a computable boundary extension even though the boundary of D is not effectively locally connected. The upshot of this is that effective local connectivity provides sufficient, but not necessary, information for computing boundary extensions of computable conformal maps. Specifically, we prove the following.

Theorem 2.1. *There is a computable and conformal map on the unit disk that has a computable boundary extension even though the image of this boundary extension on the unit circle is not effectively locally connected.*

3. An estimate on boundary behavior

Theorem 3.1. *Suppose ϕ is a conformal map on the unit disk that has a boundary extension and let D denote its range. Suppose r_0, R_0 are positive numbers such that*

$$r_0 < \frac{R_0}{\exp(16\pi^2)}$$

and such that R_0 is smaller than the minimum of $|\phi(z) - \zeta|$ as z ranges over all points whose modulus is at most $1/2$ and ζ ranges over all boundary points of D. Let ζ_0 be a boundary point of D, and choose a point $z_0 \in D$ so that ζ_0 is a boundary point of the connected component of z_0 in $D_{r_0}(\zeta_0) \cap D$. Then, $\zeta_0 = \phi(\zeta)$ for some ζ such that

$$\left|\operatorname{Arg}\left(\frac{\zeta}{\phi^{-1}(z_0)}\right)\right| < \arcsin\left(\frac{2\pi}{\sqrt{\ln(R_0/r_0)}}\right).$$

Proof. Let $z_1 = \phi^{-1}(z_0)$. Since ζ_0 is a boundary point of the connected component of z_0 in $D_{r_0}(\zeta_0) \cap D$, there is an arc β from z_0 to ζ_0 that contains no point of the boundary of D except ζ_0 and that is contained in $D_{r_0}(\zeta_0)$. Let β_1 be the pre-image of $\beta - \{\zeta_0\}$ under ϕ^{-1}.

Let

$$m = \frac{2\pi}{\sqrt{\ln(R_0/r_0)}}.$$

The upper bound on r_0 ensures that $m < 1/2$. Let ρ_1 denote the ray through 0 and z_1. Suppose ρ is a ray that is parallel to ρ_1 and whose distance from ρ_1 is m. That is, for each point z on ρ, the distance from z to ρ_1 is m. We claim that ρ does not contain any point of β_1. For, suppose otherwise. Let ρ' denote the image under ϕ of ρ. Since $m < 1/2$, it follows that ρ contains a point of $\overline{D_{1/2}(0)}$. It then follows from the upper bound on R_0 that ρ' contains a point of $D - \overline{D_{R_0}(\zeta_0)}$. But, ρ' also intersects β. It follows that ρ' intersects the circles with centers ζ_0 and radii r_0 and R_0. Let ρ_1' denote the image under ϕ of ρ_1. It similarly follows that ρ_1'

intersects theses circles. But furthermore, these images must intersect every circle whose center is ζ_0 and whose radius is between r_0 and R_0.

Let $\gamma(r,\theta) = \zeta_0 + re^{i\theta}$. Whenever $r_0 < r < R_0$, we can choose numbers $\theta_{0,r}$ and $\theta_{1,r}$ so that $\gamma(r,\theta_{0,r}) \in \rho'$, $\gamma(r,\theta_{1,r}) \in \rho'_1$, and so that $\gamma(r,\theta) \in D$ whenever θ is between $\theta_{0,r}$ and $\theta_{1,r}$. Furthermore, we can choose these numbers so that they vary continuously with r and so that $|\theta_{0,r} - \theta_{1,r}| \leq 2\pi$. Without loss of generality, suppose $\theta_{0,r_0} < \theta_{1,r_0}$. Since ρ and ρ_1 do not cross, it follows that $\theta_{0,r} < \theta_{1,r}$ whenever $r_0 < r < R_0$.

Let $\psi = \phi^{-1}$. Whenever $r_0 < r < R_0$, the modulus of $\psi\gamma(r,\theta_{0,r}) - \psi\gamma(r,\theta_{1,r})$ is at least m. At the same time, this modulus is not larger than the length of the arc traversed by $\psi\gamma(r,\theta)$ as θ ranges between $\theta_{0,r}$ and $\theta_{1,r}$. The length of this arc is

$$\int_{\theta_{0,r}}^{\theta_{1,r}} \left| \frac{\partial}{\partial\theta}\psi\gamma(r,\theta)\right| d\theta = \int_{\theta_{0,r}}^{\theta_{1,r}} |\psi'\gamma(r,\theta)| r d\theta.$$

By the Schwarz Integral Inequality,

$$\begin{aligned} m^2 &\leq \int_{\theta_{0,r}}^{\theta_{1,r}} |\psi'\gamma(r,\theta)|^2\, d\theta \int_{\theta_{0,r}}^{\theta_{1,r}} r^2\, d\theta \\ &\leq r2\pi \int_{\theta_{0,r}}^{\theta_{1,r}} |\psi'\gamma(r,\theta)|^2 r d\theta. \end{aligned}$$

Thus,

$$\frac{m^2}{2\pi}\int_{r_0}^{R_0} \frac{1}{r}\, dr \leq \int_{r_0}^{R_0}\int_{\theta_{0,r}}^{\theta_{1,r}} |\psi'\gamma(r,\theta)|^2 r d\theta dr.$$

But, since r is the Jacobian of γ, the latter integral is not larger than

$$\int_{D_{R_0}(\zeta_0)\cap D} |\psi'|^2\, dA$$

which, by the Lusin Area Integral (see, eg, Lemma 13.1.2, page 386, of Greene and Krantz [13]), is the area of a subregion of the unit disk. Hence,

$$\frac{m^2}{2\pi}\ln(R_0/r_0) \leq \pi.$$

But, this contradicts the upper bound on r_0. Thus, ρ does not intersect β_1.

So, let ρ_0, ρ_2 be the two rays that are parallel to ρ_1 and whose distance from ρ_1 is m. Let p_j denote the point where ρ_j crosses the unit circle. By the upper bound on R_0, β_1 contains no point of the circle with center 0 and radius 1/2. Let σ denote the arc on the unit circle from p_0 to p_2 that contains p_1. It follows that β_1 has a limit point on σ; let ζ denote such a point. Hence, $\phi(\zeta) = \zeta_0$. Since $\zeta \in \sigma$, it also follows that

$$\left|\mathrm{Arg}\left(\frac{\zeta}{z_1}\right)\right| < m.$$

□

4. The construction of the domain D

We begin by constructing a set $X \subseteq \mathbb{C}$. This set will be the boundary of D. To begin, we declare all points that lie on either the left, right, or top side of the unit square $[0,1]^2$ to be in X. Let $\nu_0 = 0$, $\nu_1 = i$, $\nu_2 = 1 + i$, and $\nu_3 = 1$. We then declare each point of the form $2^{-(n+1)}(1 + i)$ to be in X, and we label this point



FIGURE 1. Construction of X: the case where $n \in A$

ν_{3n+5}. (The reasons for the choice of subscript will become apparent later.) Let A be a *c.e.* but incomputable set of natural numbers, and fix an effective enumeration of A, $\{A_s\}_{s\in\mathbb{N}}$. That is, $\{A_s\}_{s\in\mathbb{N}}$ is a uniformly computable sequence of finite sets, $A_s \subseteq A_{s+1}$ for all s, and $A = \bigcup_s A_s$. We say that a number n *enters* A *at stage* s if s is the smallest number such that $n \in A_s$. If n enters A at stage s, then we declare $2^{-(n+1)} + 2^{-(n+2+s)}$ and $2^{-(n+1)} - 2^{-(n+2+s)}$ to be in X; label these points ν_{3n+4} and ν_{3n+6} respectively. See Figure 1. On the other hand, if n does not belong to A, then we only put $2^{-(n+1)}$ into X and we let $\nu_{3n+4} = \nu_{3n+6} = 2^{-(n+1)}$. See Figure 2. Then, for each n and each $j \in \{4, 6\}$, we put all points on the line segment from ν_{3n+j} to ν_{3n+5} into X. Finally, put all points in the interval $[\nu_4, 1]$ and all points in the intervals $[\nu_{3n+7}, \nu_{3n+6}]$ into X.

The set $\mathbb{C} - X$ consists of two connected components, only one of which is bounded. Let D denote the bounded component. Thus, D is simply connected and X is the boundary of D. It follows from the construction that D is computably open and X is computably closed. So, by the Effective Riemann Mapping Theorem of Hertling [16], there is a computable and conformal map ϕ of $\mathbb{D}$ onto D. The set X is locally connected, so ϕ has a boundary extension which we also denote by ϕ.

However, X is not effectively locally connected. For, suppose otherwise. Then, there is a computable map $g : \mathbb{N} \to \mathbb{N}$ with the property that whenever k is a non-negative integer and x, y are distinct points of X such that $|x - y| \leq 2^{-g(k)}$, X contains an arc from x to y whose diameter is smaller than 2^{-k}. We can assume g is increasing. Using this map g, we show that A is computable contrary to assumption. In fact, we claim that $n \notin A$ if $n \notin A_{g(n+2)}$. For, by way of contradiction, suppose that $n \notin A_{g(n+2)}$ but that $n \in A$, and let s be the stage at which n enters A. Thus, $s > g(n+2)$. It follows from the construction that the distance from ν_{3n+4} to ν_{3n+6} is $2^{-(n+1+s)}$ which in turn is smaller than $2^{-g(n+2)}$. However, by construction, the shortest arc in X from ν_{3n+4} to ν_{3n+6} consists of the line segment from ν_{3n+4} to



Figure 2. Construction of X: the case where $n \notin A$

ν_{3n+5} and the line segment from ν_{3n+5} to ν_{3n+6}. But, the diameter of each of these segments is at least $2^{-(n+1)}$, and this is a contradiction. Thus, $n \notin A$ if $n \notin A_{g(n+2)}$. However, this now contradicts the incomputability of A, and so such a function g can not exist.

5. A REPRESENTATION OF ϕ

We now construct a representation of ϕ which we will use to compute the boundary extension of ϕ. Our approach parallels that used in Andreev and McNicholl [1] to construct Schwarz-Christoffel mappings for multiply connected domains. The representation uses some of the basic elements of harmonic function theory which we summarize here.

To begin, a real-valued function u on a domain D is *harmonic* if it is twice continuously differentiable and for all $z \in D$,

$$\frac{\partial^2 u}{\partial x^2}(z) + \frac{\partial^2 u}{\partial y^2}(z) = 0.$$

The real and imaginary parts of an analytic function are harmonic. Furthermore, if u, v are the real and imaginary parts respectively of an analytic function f, then

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial v}{\partial y} &= -\frac{\partial u}{\partial x} \end{aligned}$$

These equations are called the *Cauchy-Riemann equations.* If u, v are harmonic functions on $\mathbb{D}$ that satisfy these equations, then $u+iv$ is analytic, and v is said to be a *harmonic conjugate* of u. Each harmonic function on $\mathbb{D}$ has a harmonic conjugate, and any two conjugates of a harmonic function differ by a constant. Accordingly, when u is a harmonic function on $\mathbb{D}$, we let $\tilde{u}$ be the harmonic conjugate of u with the property that $\tilde{u}(0) = 0$. When u is a harmonic function on $\mathbb{D}$ that has an integrable extension to the closure of $\mathbb{D}$, it follows from [12, Exercise II.11] that

$$\tilde{u}(re^{i\lambda}) = \int_{-\pi}^{\pi} \frac{2r\sin(\lambda - \theta)u(e^{i\theta})}{1 - 2r\cos(\lambda - \theta) + r^2}\frac{d\theta}{2\pi}. \tag{5.1}$$

We then set $\hat{u} = u + i\tilde{u}$.

If $\gamma : [a, b] \to \mathbb{C}$ is smooth, then let:

$$\begin{aligned} \int_\gamma f(\zeta)d\zeta &= \int_a^b f(\gamma(t))\gamma'(t)dt \\ \int_\gamma f(\zeta)|d\zeta| &= \int_a^b f(\gamma(t))|\gamma'(t)|dt \end{aligned}$$

If f is an integrable and real-valued function on the unit circle, and if we set

$$u(z) = \int_{\partial\mathbb{D}} f(\zeta)P_z(\zeta)|d\zeta|, \tag{5.2}$$

where

$$P_z(\zeta) = \frac{1 - |z|^2}{|\zeta - z|^2},$$

then u is a harmonic function on $\mathbb{D}$ and $f(\zeta) = \lim_{z\to\zeta} u(z)$ whenever f is continuous at ζ. The function u is called the *solution to the Dirichlét problem with boundary data f.*

If E is a Borel subset of the unit circle, then there is a unique harmonic function on $\mathbb{D}$, u, with the property that for each unimodular ζ,

$$\lim_{z\to\zeta} u(z) = \begin{cases} 1 & \zeta \in E \\ 0 & \zeta \notin E \end{cases}$$

This function is called the *harmonic measure function of E*, and its value at z is denoted $\omega(z, E, \mathbb{D})$. If we fix z and allow E to vary, then we obtain a Borel probability measure on the unit circle.

We shall refer to the points $\nu_0, \nu_1, \ldots$ as *vertices.* Following Driscoll and Trefethen [11], we then refer to their pre-images as *pre-vertices.* We then refer to all other points of the boundary of $\mathbb{D}$ as *non-prevertices.* The set of non-prevertices has infinitely many connected components.

Let J_0 be the connected component of the set of non-prevertices that is bounded by the pre-images of ν_0 and ν_1. Label the remaining components $J_1, J_2, \ldots$ so that J_n and J_{n+1} have exactly one point in common for each n.

Now, let α_0 denote the ϕ-preimage of ν_0, and for each n, let α_{n+1} denote the common point of J_n and J_{n+1}. It follows that $\phi(\alpha_n) = \nu_n$ for each n. Without loss of generality, we assume that $\alpha_0 = -1$.

Suppose J is a connected component of the set of non-prevertices. It follows from the Schwarz Reflection Principle (see p.184 of [25]) that ϕ has a differentiable extension that includes J. For all $z \in \partial\mathbb{D}$, let $\arg_1(z)$ be the number $\theta \in [-\pi, \pi)$ such that $e^{i\theta} = z$. Then, by a well-known argument, which we reproduce here for the sake of completeness, $\arg_1(z\phi'(z))$ is constant on J with respect to z. For, fix numbers θ_1 and θ_2 such that $e^{i\theta}$ belongs to J precisely when $\theta_1 < \theta < \theta_2$. Set $\eta(\theta) = \phi(e^{i\theta})$. So, by the construction of X, as θ travels from θ_1 to θ_2, $\eta(\theta)$ traces the line segment from ν_n to ν_{n+1}. Thus, as θ travels from θ_1 to θ_2, $\eta'(\theta)$

may change in magnitude but can not change in direction; that is, its argument is constant. However, by the chain rule, $\eta'(\theta) = \phi'(e^{i\theta})e^{i\theta}i$. By setting $\theta = \arg_1(z)$, we see that $\arg_1(z\phi'(z))$ is constant as z ranges over J. Moreover, if c is the value of this argument then $c = \eta'(\theta) - \pi/2$.

So, for each n let $c_n = \arg_1(z\phi'(z))$ for all $z \in J_n$. We now calculate the values of these constants. To begin, we have the following:

$$\begin{aligned} c_0 &= 0 \\ c_1 &= -\frac{\pi}{2} \\ c_2 &= -\pi \\ c_{3n} &= \frac{\pi}{2} \end{aligned}$$

When $j \notin A$, it follows that $c_{3j+4} = 0$ and that $c_{3n+5} = -\pi$. But, if j enters A at stage s, then it follows that

$$\begin{aligned} c_{3j+4} &= \arctan(2^{s+1}) \\ c_{3j+5} &= -\frac{3\pi}{2} - \arctan(2^{s+1}) \end{aligned}$$

For each $z \in \mathbb{D}$, let:

$$\begin{aligned} \Lambda(z) &= \sum_{n=0}^{\infty} c_n \omega(z, J_n, \mathbb{D}) \\ \Omega(z) &= \int_{\partial\mathbb{D}} \arg_1(\zeta) P_z(\zeta)|d\zeta| \end{aligned}$$

Let Log be the analytic branch of the logarithm whose domain is $\mathbb{C} - (-\infty, 0]$. That is,

$$\operatorname{Log}(z) = \int_1^z \frac{1}{\zeta} d\zeta.$$

Let Arg denote the imaginary part of Log. For each $z \in \partial\mathbb{D} - \{-1\}$, $\arg_1(z) = 2\operatorname{Arg}(1+z)$. The function $z \mapsto 2\operatorname{Arg}(1+z)$ is harmonic and is defined at each point in $\overline{\mathbb{D}} - \{-1\}$. It then follows that $i\hat{\Omega}(z) = 2\operatorname{Log}(1+z)$ for all $z \in \overline{\mathbb{D}} - \{-1\}$. We seek to show that

$$\phi' = C \times \exp(i\hat{\Lambda} - i\hat{\Omega}) \tag{5.3}$$

for some constant C.

To prove Equation (5.3), we begin by defining ψ to be $\exp(i\hat{\Lambda} - i\hat{\Omega})$. By the Schwarz Reflection Principle, this function has an analytic extension to an open set that contains each J_n; let ψ denote this analytic extension as well. We then consider the analytic function ψ/ϕ'. By construction, the argument of this function is 0 on each J_n. It follows that ψ/ϕ' has an analytic extension to an open set that contains $\overline{\mathbb{D}}$, and that this extension maps the boundary of the unit disk into the real line (but not the whole real line since $\partial\mathbb{D}$ is compact). So, if w is not real, then the number of times $\psi(\zeta)/\phi'(\zeta)$ travels around w as ζ travels around the unit circle is zero. Set $g = \psi/\phi'$. Then, the Winding Number

$$\frac{1}{2\pi i} \int_{g(\partial\mathbb{D})} \frac{1}{\zeta - w} d\zeta$$

is zero. But, by Change of Variables, this number is

$$\frac{1}{2\pi i}\int_{\partial\mathbb{D}}\frac{g'(\zeta)}{g(\zeta)-w}d\zeta$$

which, by the Argument Principle, is the number of pre-images of w under g that lie inside the unit disk. In other words, ψ/ϕ' maps the unit disk into the real line. By the Open Mapping Theorem, this is only possible if ψ/ϕ' is constant.

Hence, there is a constant C such that

$$\phi(z) = C\int_0^z \frac{\exp(i\hat{\Lambda}(\zeta))}{(1+\zeta)^2}d\zeta + \phi(0) \tag{5.4}$$

for all $z \in \mathbb{D}$.

6. The computability of the representation of ϕ

Our goal is to use the representation (5.4) to demonstrate the computability of the boundary extension of ϕ. To do so, we first must compute the value of C. Since ϕ' is computable, (see eg Proposition 4.1 of [16]), this amounts to showing that $\hat{\Lambda}$ and $\hat{\Omega}$ are computable. By (5.1), it suffices to show that Λ and Ω are computable. The computability of Ω is obvious. In order to demonstrate the computability of Λ, we first note that the sequence $\{c_j\}_{j\in\mathbb{N}}$ is computable. We then seek to show that $\omega(z, J_n, \mathbb{D})$ is a computable function of n, z which reduces to showing that $\{\alpha_n\}_{n\in\mathbb{N}}$ is computable. So, let (n, k) be given as input. We seek to compute a rational point q so that $|q - \alpha_n| < 2^{-k}$. We apply Theorem 3.1 with ν_n playing the role of ζ_0. In finitely many steps, we can compute a rational number R_0 such that

$$0 < R_0 < \min\{|\phi(z) - \zeta| \mid |z| \leq \frac{1}{2}, \zeta \in \partial D\}.$$

Compute a rational number $r_0 < R_0/\exp(16\pi^2)$ so that

$$1 - \sqrt{1-m^2} \quad < \quad 2^{-(k+2)} \tag{6.1}$$

where

$$m = \frac{2\pi}{\sqrt{\log(R_0/r_0)}}.$$

We now compute a point $z_0 \in D \cap D_{r_0}(\zeta_0)$ such that ζ_0 is a boundary point of the connected component of z_0 in $D_{r_0}(\zeta_0) \cap D$. To do so requires that we take into consideration the value of n. If $n = 0$, then we set $z_0 = \frac{r_0}{2}(1+2i)$ so that z_0 lies on the line $y = 2x$. If $n = 1$, then we set $z_0 = \frac{r_0}{2} + i(1 - \frac{r_0}{2})$ so that $|z_0 - \nu_1| = \frac{r_0}{\sqrt{2}}$ and z_0 lies on the line $y = 1 - x$. If $n = 2$, then we set $z_0 = (1 - \frac{r_0}{2})(1 + i)$ so that $|z_0 - \nu_2| = r_0/\sqrt{2}$ and z_0 lies on the line $y = x$. If $n = 3$, then let $z_0 = (1 - \frac{r_0}{3}) + 2(\frac{r_0}{3})i$ so that

$$|z_0 - \nu_3| = \left|\frac{-r_0}{3} + \frac{2r_0}{3}i\right| = \frac{\sqrt{5}}{3}r_0$$

and so that z_0 lies on the line $y = -2(x-1)$.

If $n = 3j + 5$, then set $z_0 = \nu_n + \frac{r_0}{2}i$. Suppose $n = 3j + b$ where $b \in \{4, 6\}$. Compute the smallest s such that $j \in A_s$ or $2^{-(n+s+2)} < \frac{r_0}{2}$. Set

$$z_0 = 2^{-(n+1)} \pm 2^{-(n+s+2)} + \frac{r_0}{2}i$$

where the sign is positive if $b = 4$ and negative if $b = 6$. If $j \in A_s$, then $z_0 = \nu_n + \frac{r_0}{2} i$. If $j \notin A_s$, then by the choice of s, $|z_0 - \nu_n| < r_0$.

In all of these cases, it follows that every point between ν_n and z_0 lies in D.

Let $\rho = |\phi^{-1}(z_0)|$, and let $w_0 = \rho^{-1}\phi^{-1}(z_0)$. It follows from Theorem 3.1 that there is a pre-image of ν_n, ξ, such that

$$|\xi - w_0| \leq 2 - 2\sqrt{1 - m^2} < 2^{-(k+1)}.$$

Compute a rational point q such that $|q - w_0| < 2^{-(k+1)}$. Thus, $|q - \xi| < 2^{-k}$. If n is 0, 1, 2, or 3, or if n is congruent to 5 modulo 3, then ν_n has only one pre-image and so $|q - \alpha_n| < 2^{-k}$. Suppose $n = 3j + 6$ where $j \in \mathbb{N}$. Then, ν_n may have two pre-images under ϕ. That is, if $j \in A$, then $\nu_n = \nu_{n-2}$ and $\phi(\alpha_n) = \phi(\alpha_{n-2}) = \nu_n$. However, α_n lies in the subarc of $\partial\mathbb{D}$ from α_{n-1} to α_{n+2} that does not contain α_{n+5}, and α_{n-2} does not belong to this arc. This arc can now be computed. So, we choose r_0 small enough so that this subarc contains

$$\{\zeta \in \partial\mathbb{D} \ : \ |\operatorname{Arg}(\zeta/w_0)| \leq m\}.$$

So, by 6.1, $|w_0 - \alpha_n| < 2^{-(k+1)}$. Hence, $|q - \alpha_n| < 2^{-k}$. The remaining cases are handled similarly.

We have now shown that Λ is computable. It now follows that $\hat{\Lambda}$ is computable. As discussed in the introduction to this section, it follows that C is computable. We are now ready to demonstrate that the boundary extension of ϕ is computable.

7. A PROOF OF THE COMPUTABILITY OF THE BOUNDARY EXTENSION OF ϕ

It will be useful later in this section to have a computable parametrization of X. This can be constructed by first building, for each $t \in \mathbb{N}$, a stage t approximation of X, X_t, by making the following modifications to the construction of X. If n enters A at stage $s \leq t$, then we declare $2^{-(n+1)} + 2^{-(n+2+s)}$ and $2^{-(n+1)} - 2^{-(n+2+s)}$ to be in X_t. On the other hand, if n does not belong to A_t, then put $2^{-(n+1)} + 2^{-(n+2+t)}$ and $2^{-(n+1)} - 2^{-(n+2+t)}$ into X_t. The rest of the construction of X is unchanged. The resulting set X_t is a Jordan curve with a computable parameterization γ_t. We can compute γ_t so that $\gamma_t(s_1) = \gamma_t(s_2)$ if and only if $s_1, s_2 \in \{0, 1\}$. We can further choose $\gamma_0, \gamma_1, \ldots$ so that they converge uniformly to a surjection γ of $[0,1]$ onto X. Furthermore, we can ensure that from k one can compute a number t_0 so that $|\gamma_{t_1}(s) - \gamma_{t_2}(s)| < 2^{-k}$ whenever $t_1, t_2 \geq t_0$ and $s \in [0,1]$. Hence, γ is computable. We note that if $n \notin A$, then $2^{-(n+1)}$ and each point between $2^{-(n+1)}$ and $2^{-(n+1)}(1 + i)$ has exactly two γ-preimages; all other points on X have exactly one γ-preimage.

The rest of the proof now turns on the following lemma which is a consequence of [22, Lemma 2] which roughly states that Dirichlét problems and their boundary extensions are uniformly computable from piecewise continuous data.

Lemma 7.1. *If f is a computable function on the unit circle, and if u is the solution to the resulting Dirichlét problem, then the continuous extension of u to the closed unit disk is computable.*

We now show that the boundary extension of ϕ is computable. To this end, we first show that we can compute ϕ on the unit circle. For, we can then apply Lemma 7.1 to compute the real and imaginary parts of the boundary extension of ϕ.

As a step in this direction, suppose θ is a rational number and that $e^{i\theta}$ is neither a pre-vertex nor -1. We show that $\phi(e^{i\theta})$ is computable uniformly in θ. That is, given

θ, k as input, one can compute a rational point q such that $|q - \phi(e^{i\theta})| < 2^{-k}$. For the moment, set $w = e^{i\theta}$ and fix rational numbers r_1, r_2 such that $0 < r_1 < r_2 < 1$. Then,

$$\begin{aligned} |\phi(r_1 w) - \phi(r_2 w)| &= |C| \left| \int_{r_1 w}^{r_2 w} \frac{\exp(i\hat{\Lambda}(\zeta))}{(1+\zeta)^2} d\zeta \right| \\ &\leq |C| \int_{r_1 w}^{r_2 w} \frac{|\exp(i\hat{\Lambda}(\zeta))|}{|1+\zeta^2|} |d\zeta| \\ &\leq |C| \frac{M_1}{M_2^2} |r_2 - r_1| \end{aligned}$$

where M_1 is the supremum of $|\exp(i\hat{\Lambda}(rw))|$ as r ranges from r_1 to r_2, and M_2 is the infimum of $|1 + rw|$ as r ranges from r_1 to r_2. M_2 can be computed using basic calculus. To estimate M_1, we first observe that $|\exp(i\hat{\Lambda}(z))| = |\exp(-\tilde{\Lambda}(z))|$. We then observe that by (5.1), if $z = re^{i\lambda}$, then

$$\tilde{\Lambda}(z) = \int_{-\pi}^{\pi} \frac{2r\sin(\lambda - \theta)\Lambda(e^{i\theta})}{1 - 2r\cos(\lambda - \theta) + r^2} \frac{d\theta}{2\pi}.$$

We therefore seek a lower bound on this integral. We begin by noticing that

$$\frac{2r\sin(\lambda - \theta)}{1 - 2r\cos(\lambda - \theta) + r^2} \geq \frac{-2r}{(1+r)^2}.$$

Since $\omega(z, E, \mathbb{D})$, as a function of E, is a probability measure on the boundary of E, $\sum_{n=0}^{\infty} \omega(z, J_n, \mathbb{D}) = \omega(z, \partial\mathbb{D}, \mathbb{D}) = 1$. As $-\pi \leq c_n \leq \pi$, we can conclude that $-\pi \leq \Lambda(e^{i\theta}) \leq \pi$. Hence,

$$\tilde{\Lambda}(z) \geq -\frac{2\pi r}{(1+r)^2} \geq -\frac{\pi}{2}.$$

This enables us to infer that $M_1 \leq \exp(\pi/2)$. By allowing r_2 to approach 1 from the left, we conclude that $\phi(r_1 e^{i\theta})$ can be used to estimate $\phi(e^{i\theta})$ with error at most $\exp(\pi/2)(1 - r_1)/M_2^2$. Since $\phi(r_1 e^{i\theta})$ can be computed uniformly in r_1 and θ, it follows that $\phi(e^{i\theta})$ can be computed uniformly in θ.

We now show that ϕ can be computed on the unit circle. Suppose we are given as input an open rational interval (θ_1, θ_2). It is required to produce an open rational disk that contains $\phi(e^{i\theta})$ whenever $\theta_1 < \theta < \theta_2$. We begin by simultaneously computing each $e^{i\theta_j}$ and each α_n until we obtain good enough approximations to enough of these quantities to demonstrate that neither $e^{i\theta_1}$ nor $e^{i\theta_2}$ is a pre-vertex or -1 and to also demonstrate that for some n_0, $\alpha_{n_0} \neq e^{i\theta}$ whenever $\theta_1 \leq \theta \leq \theta_2$. If no such stage is found, then this computation will not halt. However, for each real θ there are arbitrarily close rational numbers θ_1 and θ_2 such that $\theta_1 < \theta < \theta_2$ and this search terminates.

We then continue by computing each $\phi(e^{i\theta_j})$. Once these have been approximated with sufficient closeness, we can begin computing t_1 and t_2 so that $\gamma(t_j) = \phi(e^{i\theta_j})$ and $\alpha_{n_0} \neq \gamma(t)$ whenever t is between t_1 and t_2. It follows that whenever $\theta_1 < \theta < \theta_2$, $\phi(e^{i\theta}) = \gamma(t)$ for some t between t_1 and t_2. Let g be an increasing and computable modulus of continuity for γ. Calculate the smallest k so that $|t_1 - t_2| \leq 2^{-g(k)}$. Compute a rational point q so that $|q - \gamma(t_1)| < 2^{-k}$. Then, $|q - \phi(e^{i\theta})| < 2^{-k+1}$ whenever $\theta_1 < \theta < \theta_2$. The quantity $|t_2 - t_1|$ approaches zero as $\theta_2 - \theta_1$ approaches zero. Thus the Convergence criterion is satisfied (see Section 2). It follows that ϕ is computable on the unit circle and hence on all of $\overline{\mathbb{D}}$.

8. Conclusion

9. Acknowledgement

My deepest thanks to the referees for their many helpful comments and to Justin Peters for helpful conversations.

References

1. V.V. Andreev and T.H. McNicholl, *A potential-theoretic construction of the Schwarz-Christoffel map for finitely connected domains*, Complex Variables and Elliptic Equations **58** (2013), no. 2, 163 – 183.
2. I. Binder, M. Braverman, and M. Yampolsky, *On the computational complexity of the Riemann mapping*, Archiv for Matematik **45** (2007), 221–239.
3. Errett Bishop and Douglas Bridges, *Constructive analysis*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 279, Springer-Verlag, Berlin, 1985.
4. V. Brattka, *Effective borel measurability and reducibility of functions*, Mathematical Logic Quarterly **51** (2005), no. 1, 19 – 44.
5. Vasco Brattka and Klaus Weihrauch, *Computability on subsets of Euclidean space. I. Closed and compact subsets*, Theoret. Comput. Sci. **219** (1999), no. 1-2, 65–93, Computability and complexity in analysis (Castle Dagstuhl, 1997).
6. H. Cheng, *A constructive Riemann mapping theorem*, Pacific Journal Mathematics **44** (1973), 435 – 454.
7. S. Barry Cooper, *Computability theory*, Chapman & Hall/CRC, Boca Raton, FL, 2004.
8. P.J. Couch, B.D. Daniel, and T.H. McNicholl, *Computing space-filling curves*, Theory of Computing Systems **50** (2012), no. 2, 370–386.
9. Nigel Cutland, *Computability*, Cambridge University Press, Cambridge, 1980, An introduction to recursive function theory.
10. D. Daniel and T.H. McNicholl, *Effective local connectivity properties*, Theory of Computing Systems **50** (2012), no. 4, 621 – 640.
11. Tobin A. Driscoll and Lloyd N. Trefethen, *Schwarz-Christoffel mapping*, Cambridge Monographs on Applied and Computational Mathematics, vol. 8, Cambridge University Press, Cambridge, 2002.
12. J. B. Garnett and D. E. Marshall, *Harmonic measure*, New Mathematical Monographs, vol. 2, Cambridge University Press, Cambridge, 2005.
13. R. Greene and S. Krantz, *Function theory of one complex variable*, Graduate Studies in Mathematics, American Mathematical Society, 2002.
14. X. Gu, J.H. Lutz, and E. Mayordomo, *Curves that must be retraced*, Information and Computation **209** (2011), 992 – 1006.
15. P. Henrici, *Applied and computational complex analysis. Vol. 3*, Pure and Applied Mathematics (New York), John Wiley & Sons Inc., New York, 1986, Discrete Fourier analysis—Cauchy integrals—construction of conformal maps—univalent functions, A Wiley-Interscience Publication.
16. P. Hertling, *An effective Riemann Mapping Theorem*, Theoretical Computer Science **219** (1999), 225 – 265.
17. Iraj Kalantari and Lawrence Welch, *Point-free topological spaces, functions and recursive points; filter foundation for recursive analysis. I*, Ann. Pure Appl. Logic **93** (1998), no. 1-3, 125–151.
18. Akitoshi Kawamura and Stephen Cook, *Complexity theory for operators in analysis*, STOC'10—Proceedings of the 2010 ACM International Symposium on Theory of Computing (New York), ACM, 2010, pp. 495–502.
19. Ker-I Ko, *Complexity theory of real functions*, Progress in Theoretical Computer Science, Birkhäuser, Boston, 1991.

20. P. Koebe, *Über eine neue Methode der Konformen Abbildung und Uniformisierung*, Nachr. Kgl. Ges. Wiss. Göttingen, Math.-Phys. Kl. 1912 (1912), 844–848.
21. T.H. McNicholl, *Computing boundary extensions of conformal maps*, Submitted. Preprint available at http://arxiv.org/abs/1110.5271v1.
22. ______, *An effective Carathéodory theorem*, Theory of Computing Systems **50** (2012), no. 4, 579 – 588.
23. ______, *Computing links and accessing arcs*, Mathematical Logic Quarterly **59** (2013), no. 1 - 2, 101 – 107.
24. J. Miller, *Effectiveness for embedded spheres and balls*, CCA 2002, Computability and Complexity in Analysis (V. Brattka, M. Schröder, and K. Weihrauch, eds.), vol. 66, Electronic Notes in Computer Science, no. 1, Elsevier, 2002, pp. 127 – 138.
25. Zeev Nehari, *Conformal mapping*, McGraw-Hill Book Co., Inc., New York, Toronto, London, 1952.
26. Ch. Pommerenke, *Boundary behaviour of conformal maps*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 299, Springer-Verlag, Berlin, 1992.
27. Marian B. Pour-El and J. Ian Richards, *Computability in analysis and physics*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1989.
28. Klaus Weihrauch, *Computable analysis*, Texts in Theoretical Computer Science. An EATCS Series, Springer-Verlag, Berlin, 2000.

Department of Mathematics, Iowa State University, Ames, Iowa 50011 USA
E-mail address: `mcnichol@iastate.edu`